


## CHEMISTRY-CY

## Q.1 – Q.25 : Carry ONE mark each.

1. Amongst the following the compound that **DOES NOT** act as a diene in Diels-Alder reaction is



2. An efficient catalyst for hydrogenation of alkenes is  $[\text{Rh}(\text{PPh}_3)_3\text{Cl}]$ . However,  $[\text{Ir}(\text{PPh}_3)_3\text{Cl}]$  does not catalyze this reaction, because

- (a)  $\text{PPh}_3$  binds stronger to Ir than to Rh (b) Cl binds stronger to Ir than to Rh  
(c)  $\text{PPh}_3$  binds stronger to Rh than to Ir (d) Cl binds stronger to Rh than to Ir


3. Which of the following properties are characteristics of an ideal solution?

- (i)  $(\Delta_{\text{mix}}G)_{T,P}$  is negative (ii)  $(\Delta_{\text{mix}}S)_{T,P}$  is positive  
(iii)  $(\Delta_{\text{mix}}V)_{T,P}$  is positive (iv)  $(\Delta_{\text{mix}}H)_{T,P}$  is negative.  
(a) (i) and (iv) (b) (i) and (ii) (c) (i) and (iii) (d) (iii) and (iv)

4. Among the following compounds, the one that is non-aromatic, is




5. Given the  $E^0$  values for the following reaction sequence,



the computed value of  $E^0$  for  $\text{Mn}^{6+} \rightarrow \text{Mn}^{2+}$  (in volts) is \_\_\_\_\_

6. The expression for the equilibrium constant ( $K_{\text{eq}}$ ) for the enzyme catalyzed reaction given below, is




- (a)  $\frac{k_1 k_3}{k_2 k_4}$  (b)  $\frac{k_1 k_2}{k_3 k_4}$  (c)  $\frac{k_2 k_3}{k_1 k_4}$  (d)  $\frac{k_1 k_4}{k_2 k_3}$

7. The absorption spectrum of  $[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$  in solution comprises of a maximum with a shoulder. The reason for the shoulder is


- (a) ligand-to-metal charge transfer (LMCT) (b) metal-to-ligand charge transfer (MLCT)  
(c) Jahn-Teller distortion (d) nephelauxetic effect.

8. The compound given below is a




- (a) sesterterpene (b) monoterpene (c) sesquiterpene (d) triterpene

9. The electrical conductivity of a metal  
 (a) increases with increasing temperature (b) decreases with increasing temperature  
 (c) is independent of temperature (d) shows oscillatory behaviour with temperature
10. Which one of the following statements is **INCORRECT**?  
 (a) Frenkel defect is a cation vacancy and a cation interstitial  
 (b) Frenkel defect is an anion vacancy and a cation interstitial  
 (c) Density of a solid remains unchanged in case of Frenkel defects.  
 (d) Density of a solid decreases in case of Schottky defects.
11. Among the given pH values, the  $O_2$  binding efficiency of hemoglobin is maximum at  
 (a) 6.8 (b) 7.0 (c) 7.2 (d) 7.4
12. When the operator,  $-\hbar^2 d^2/dx^2$ , operates on the function  $e^{-ikx}$ , the result is  
 (a)  $k^2 \hbar^2 e^{-ikx}$  (b)  $ik^2 \hbar^2 e^{-ikx}$  (c)  $i \hbar^2 e^{-ikx}$  (d)  $\hbar^2 e^{-ikx}$




From the above Carnot cycle undergone by an ideal gas, identify the processes in which the change in internal energy is **NON-ZERO**.

- (a) I and II (b) II and IV (c) II and III (d) I and IV
14. Which one of the following defines the absolute temperature of a system?  
 (a)  $\left(\frac{\partial U}{\partial S}\right)_V$  (b)  $\left(\frac{\partial A}{\partial S}\right)_V$  (c)  $\left(\frac{\partial H}{\partial S}\right)_V$  (d)  $\left(\frac{\partial G}{\partial S}\right)_V$
15. The following conversion is an example of




- (a) Arndt-Eistert homologation (b) Mannich reaction  
 (c) Michael addition (d) Chichibabin amination reaction
16. The compound with planar geometry is  
 (a)  $N(t\text{-Bu})_3$  (b)  $NPh_3$  (c)  $NF_3$  (d)  $N(SiH_3)_3$
17. Reaction of benzaldehyde and p-methylbenzaldehyde under McMurry coupling conditions ( $TiCl_3$  and  $LiAlH_4$ ) gives a mixture of alkenes. The number of alkenes formed is \_\_\_\_\_
18. The correct order of reactivity of p-halonitrobenzenes in the following reaction is




- (a) p-chloronitrobenzene > p-iodonitrobenzene > p-fluoronitrobenzene > p-bromonitrobenzene  
 (b) p-fluoronitrobenzene > p-chloronitrobenzene > p-bromonitrobenzene > p-iodonitrobenzene  
 (c) p-iodonitrobenzene > p-bromonitrobenzene > p-chloronitrobenzene > p-fluoronitrobenzene  
 (d) p-bromonitrobenzene > p-fluoronitrobenzene > p-iodonitrobenzene > p-chloronitrobenzene


19. The absolute configuration of C2 and C3 in the following compound is



- (a) 2R, 3S                      (b) 2S, 3R                      (c) 2S, 3S                      (d) 2R, 3R
20. For an ideal gas with molar mass  $M$ , the molar translational entropy at a given temperature is proportional to  
 (a)  $M^{3/2}$                       (b)  $M^{1/2}$                       (c)  $e^M$                       (d)  $\ln(M)$
21. Tollen's test is **NEGATIVE** for  
 (a) mannose                      (b) maltose                      (c) glucose                      (d) sucrose
22. The intense red color of  $[\text{Fe}(\text{bpy})_3]^{2+}$  (bpy = 2, 2'-bipyridine) is due to  
 (a) metal-to-ligand charge transfer (MLCT)                      (b) ligand-to-metal charge transfer (LMCT)  
 (c) d-d transition                      (d) inter-valence charge transfer (IVCT)
23. The ease of formation of the adduct,  $\text{NH}_3 \cdot \text{BX}_3$  (where  $\text{X} = \text{F}, \text{Cl}, \text{Br}$ ) follows the order  
 (a)  $\text{BBr}_3 < \text{BCl}_3 < \text{BF}_3$                       (b)  $\text{BCl}_3 < \text{BF}_3 < \text{BBr}_3$   
 (c)  $\text{BF}_3 < \text{BCl}_3 < \text{BBr}_3$                       (d)  $\text{BBr}_3 < \text{BF}_3 < \text{BCl}_3$
24. Which one of the following plots represents an acceptable wavefunction?




25. The mass spectrum of a dihalo compound shows peaks with relative intensities of 1 : 2 : 1 corresponding to  $M$ ,  $M + 2$  and  $M + 4$  ( $M$  is the mass of the molecular ion), respectively. The compound is




**Q.26 – Q.55 : Carry TWO marks each.**

26. The value of 'g' and the number of signals observed for the reference standard, diphenylpicrylhydrazyl (DPPH), in the solid state ESR spectrum are, respectively  
(a) 2.0036 and 1      (b) 2.0036 and 3      (c) 2.2416 and 1      (d) 2.2416 and 3
27. Solvolysis of the optically active compound X gives, mainly




28. The complexes  $K_2[NiF_6]$  and  $K_3[CoF_6]$  are  
(a) both paramagnetic  
(b) both diamagnetic  
(c) paramagnetic and diamagnetic, respectively  
(d) diamagnetic and paramagnetic, respectively
29. The tetrapeptide, Ala-Val-Phe-Met, on reaction with Sanger's reagent, followed by hydrolysis gives




30. For a gas phase unimolecular reaction at temperature 298K, with a pre-exponential factor of  $2.17 \times 10^{13} \text{ s}^{-1}$ , the entropy of activation ( $\text{JK}^{-1} \text{ mol}^{-1}$ ) is \_\_\_\_\_

31. The process given below follows the Langmuir adsorption isotherm




If  $\theta$  denotes the surface coverage and  $P$  denotes the pressure, the slope of the plot of  $1/\theta$  versus  $1/\sqrt{P}$  is

- (a)  $1/(K_{eq})^2$       (b)  $1/K_{eq}$       (c)  $-1/K_{eq}$       (d)  $1/(K_{eq})^{1/2}$
32. The major product formed in the following reaction is




33. A liquid has vapor pressure of  $2.02 \times 10^3 \text{ N m}^{-2}$  at 293 K and heat of vaporization of  $41 \text{ kJ mol}^{-1}$ . The boiling point of the liquid (in Kelvin) is \_\_\_\_\_
34. The difference in the ground state energies (kJ/mol) of an electron in one-dimensional boxes of lengths 0.2 nm and 2 nm is \_\_\_\_\_
35. The major product formed in the following reaction is







36. The internal energy of an ideal gas follows the equation  $U = 3.5 PV + k$ , where  $k$  is a constant. The gas expands from an initial volume of  $0.25 \text{ m}^3$  to a final volume of  $0.86 \text{ m}^3$ . If the initial pressure is  $5 \text{ N m}^{-2}$ , the change in internal energy (in joules) is (given  $PV^{1/3} = \text{constant}$ ) \_\_\_\_\_
37. One mole of a substance is heated from  $300\text{K}$  to  $400\text{K}$  at constant pressure. The  $C_p$  of the substance is given by,  $C_p (\text{JK}^{-1}\text{mol}^{-1}) = 5 + 0.1T$ . The change in entropy, in  $\text{JK}^{-1} \text{mol}^{-1}$ , of the substance is \_\_\_\_\_
38. The solubility product of  $\text{AgBr(s)}$  is  $5 \times 10^{-13}$  at  $298\text{K}$ . If the standard reduction potential of the half-cell,  $E_{\text{Ag}|\text{AgBr(s)}|\text{Br}^-}^0$  is  $0.07\text{V}$ , the standard reduction potential,  $E_{\text{Ag}^+|\text{Ag}}^0$  (in volts) is \_\_\_\_\_
39. The most suitable reagent (s) to effect the following transformation is




- (a)  $\text{N}_2\text{H}_4$ ,  $\text{KOH}$  heat  
(c)  $\text{LiAlH}_4$
- (b)  $\text{TsNHNH}_2$ ,  $\text{CF}_3\text{COOH}$   
(d)  $\text{Na}$ , liq.  $\text{NH}_3$
40. Ammonolysis of  $\text{S}_2\text{Cl}_2$  in an inert solvent gives  
(a)  $\text{S}_2\text{N}_2$  (b)  $\text{S}_2\text{N}_2\text{Cl}$  (c)  $\text{S}_2\text{N}_2\text{H}_4$  (d)  $\text{S}_4\text{N}_4$
41. The mean ionic activity coefficient of  $0.001 \text{ molal ZnSO}_4 (\text{aq})$  at  $298\text{K}$  according to the Debye-Huckel limiting law is (Debye-Huckel constant is  $0.509 \text{ molal}^{-1/2}$ ) \_\_\_\_\_
42. Identify the function of hemocyanin and the metal responsible for it  
(a)  $\text{O}_2$  transport and  $\text{Fe}$  (b)  $\text{O}_2$  transport and  $\text{Cu}$   
(c) electron transport and  $\text{Fe}$  (d) electron transport and  $\text{Cu}$
43. The point group of  $\text{IF}_7$  is  
(a)  $D_{6h}$  (b)  $D_{5h}$  (c)  $C_{6v}$  (d)  $C_{5v}$
44. The limiting current (in  $\mu\text{A}$ ) from the reduction of  $3 \times 10^{-4} \text{ M Pb}^{2+}$ , using a dropping mercury electrode (DMF) with characteristics,  $m = 3.0 \text{ mg s}^{-1}$  and  $t = 3\text{s}$ , is (diffusion coefficient of  $\text{Pb}^{2+} = 1.2 \times 10^{-5} \text{ cm}^2\text{s}^{-1}$ ) \_\_\_\_\_
45. Identify  $X$  in the reaction,  $[\text{Pt}(\text{NH}_3)_4]^{2+} + 2\text{HCl} \rightarrow X$   
(a)  $\text{cis} - [\text{PtCl}_2(\text{NH}_3)_2]$  (b)  $\text{trans} - [\text{PtCl}_2(\text{NH}_3)_2]$   
(c)  $[\text{PtCl}(\text{NH}_3)_3]^+$  (d)  $[\text{PtCl}_3(\text{NH}_3)]^-$
46. The major products,  $K$  and  $L$  formed in the following reactions are








47. The major product formed in the following reaction is




48. The percent transmittance of  $8 \times 10^{-5}$  M solution of  $\text{KMnO}_4$  is 39.8 when measured at 510 nm in a cell of path length of 1 cm. The absorbance and the molar extinction coefficient (in  $\text{M}^{-1} \text{cm}^{-1}$ ) of this solution are, respectively  
 (a) 0.30 and 4500 (b) 0.35 and 4800 (c) 0.4 and 5000 (d) 0.48 and 5200
49. The rotational partition function of a diatomic molecule with energy levels corresponding to  $J = 0, 1$ , is (where,  $\epsilon$  is a constant)  
 (a)  $1 + e^{-2\epsilon}$  (b)  $1 + 3e^{-2\epsilon}$  (c)  $1 + e^{-3\epsilon}$  (d)  $1 + 3e^{-3\epsilon}$
50. When one CO group is replaced by  $\text{PPh}_3$  in  $[\text{Cr}(\text{CO})_6]$ , which one of the following statement is TRUE?  
 (a) The Cr-C bond length increases and CO bond length decreases  
 (b) The Cr-C bond length decreases and CO bond length decreases  
 (c) The Cr-C bond length decreases and CO bond length increases  
 (d) The Cr-C bond length increases and CO bond length increases
51. The number of possible stereoisomers obtained in the following reaction is \_\_\_\_\_



52. The major product formed in the following reaction is




53. The Beckmann rearrangement of a bromoacetophenone oxime ( $\text{C}_8\text{H}_8\text{BrNO}$ ) gives a major product having the following  $^1\text{H}$  NMR ( $\delta$ , ppm) : 9.89 (s, 1H), 7.88 (s, 1H), 7.45 (d, 1H,  $J = 7.2$  Hz), 7.17 (m, 1H), 7.12 (d, 1H,  $J = 7.0$  Hz), 2.06 (s, 3H). The structure of the product is



54. The distance between two successive (110) planes in a simple cubic lattice with lattice parameter 'a' is



55. The potential energy (PE) versus reaction coordinate diagrams for electron transfer reactions with rate constants  $k_1$ ,  $k_2$  and  $k_3$ , are given below. The increasing order of the rate constants is



Reaction coordinate

- (a)  $k_2 < k_3 < k_1$  (b)  $k_2 < k_1 < k_3$  (c)  $k_3 < k_2 < k_1$  (d)  $k_3 < k_1 < k_2$

\*\*\*\*\* END OF THE QUESTION PAPER \*\*\*\*\*